

**Use Germ-A-Cide 64 in:** Restaurants, Schools, Bathrooms, Dressing Rooms, Locker Rooms, Shower and Bath Areas, Exercise Facilities, Kennels, Pet Shops, Zoos. Use Germ-A-Cide 64 on washable hard, nonporous surfaces of: Countertops, Bathroom Fixtures, Floors. Germ-A-Cide 64 is effective at eliminating odors on porous surfaces such as drapes, carpets, bedding, shower curtains and non-porous surfaces such as walls and floors. Excellent for fire restoration.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DISINFECTION	
DILUTION:	1:64
660 ppm quat	
	2 ounces per gallon of water
	4 ounces per 2 gallons of water
	8 ounces per 4 gallons of water
	10 ounces per 5 gallons of water
	12 ounces per 6 gallons of water

**DIRECTIONS FOR USE**  
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
This product is not for use on medical device surfaces.

**DISINFECTING/CLEANING/DEODORIZING DIRECTIONS:** Remove visible soil deposits from surface, then visibly wet surface with a use-solution of 2 ounces of the concentrate per gallon of water. Use 8 oz per gallon of water to kill Adenovirus Type 7. The use-solution can be applied with a cloth, mop, sponge, or coarse spray or by soaking. For sprayer applications, use a coarse spray device. Spray 6-8 inches from the surface, rub with a brush, cloth or sponge. Do not breathe spray. Let solution remain visibly wet on surface for a minimum of 10 minutes. Rinse or allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished.

**For SARS-CoV-2, treated surfaces\*, must remain visibly wet for 1 minute.**  
**For contact surfaces\***, must be thoroughly rinsed with potable water. This product must not be used to clean the following food contact surfaces: utensils, glassware and dishes.

**For Human Coronavirus**, treated surfaces must remain visibly wet for 1 minute. Wipe dry with a clean cloth, sponge or mop or allow to air dry.  
**For Influenza Virus Type A**, treated surfaces must remain visibly wet for 2 minutes. Wipe dry with a clean cloth, sponge or mop or allow to air dry.

Prepare a fresh use-solution daily or more often if the solution becomes visibly dirty or diluted.  
**FUNGICIDAL DIRECTIONS:** For use in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities follow disinfection directions.

**MILDEWSTATIC INSTRUCTIONS:** Will effectively control the growth of mold and mildew, plus the odors caused by them when applied to hard, nonporous surfaces such as walls, floors and table tops. Apply use-solution of 2 ounces per gallon of water with a cloth, mop, sponge or coarse spray. Make sure to visibly wet all surfaces completely. Let air dry. Repeat application weekly or when growth reappears.

**\*KILLS HIV-1(AIDS VIRUS) AND HBV(HEPATITIS B VIRUS) AND HCV(HEPATITIS C VIRUS) ON PRECLEANED, HARD NON-POROUS ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood / body fluids, and in which the surfaces / objects likely to be soiled with blood / body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type (HIV-1) or Hepatitis B virus(HBV) or Hepatitis C virus(HCV).

**Special Instructions for Disinfection, Cleaning and Decontamination against HIV-1 or HBV or HCV on Surfaces (Objects Soiled with Blood / Body Fluids):**  
**Personal Protection:** Disposable protective gloves, gowns, face masks and eye coverings, must be worn during all cleaning of blood / body fluids and during decontamination procedures.

**Cleaning Procedures:** Blood / body fluids must be visibly cleaned from surfaces / objects before application of disinfectant.  
**Contact Time:** Mix 2 oz. of Germ-A-Cide 64 per gallon of water; HIV-1 is inactivated after a contact time of 4 minutes at 25°C. HBV and HCV are inactivated after a 10 minute contact time. Use appropriate contact time as listed on label for all other viruses, bacteria and fungi claimed.

**Disposal of Infectious Materials:** Blood / body fluids must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.  
**VETERINARY PRACTICE / ANIMAL CARE / ANIMAL LABORATORY / ZOO'S / PET SHOP / KENNELS DISINFECTION DIRECTIONS:**

All cleaning and disinfecting hard nonporous surfaces: equipment used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Visibly clean all surfaces with soap and detergent and rinse with water. Saturate surfaces with a use-solution of 2 oz. of Germ-A-Cide 64 per gallon of water. Allow surfaces to remain visibly wet for a period of 10 minutes. Use 8 oz. per gallon of water to kill Adenovirus Type 7. Immerse all animal handling and restraining equipment as well as forks, shovels and scrapers used to remove litter and manure. Wipe or allow to air dry. Ventilate buildings and other closed spaces. Thoroughly scrub all treated surfaces that come into contact with food, including equipment used for feeding or watering, with soap or detergent and rinse with potable water before reuse. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

**FOR HEAVY DUTY CLEANING AND / OR ODOR CONTROL:** When greater cleaning is desired, use 4 - 8 oz. of Germ-A-Cide 64 per gallon of water. (Visibly soiled areas may require repeated cleaning before treatment.)

**Deodorizing Directions:**  
Carpet: 2 oz. per gallon of shampoo solution eliminates odors associated with urine, vomit, smoke, and mildew. Reapply after cleaning for freshening effects. Allow to dry.

**Sewage Backup, Water, Smoke Damage:** Use 2 - 4 oz. per gallon of water. Spray over affected areas before and after cleaning and extraction. Allow to remain visibly wet for a contact time of ten minutes. Use proper ventilation; open windows.

APPROVED  
By: C.Reyes-Garcia at 11:51 am, Dec 13, 2021

# Germ-A-Cide 64

Per EPA stamped label 04/06/2021 ONE-STEP DISINFECTANT

## GERMICIDAL DETERGENT AND DEODORANT

- Disinfectant - Pseudomonacidal<sup>1</sup> - Staphylocidal<sup>1</sup> -
- Salmonellacidal<sup>4</sup> - Bactericidal - Fungicidal -
- Mildewstatic - \*Virucidal

**Active Ingredients:**

Didecyl dimethyl ammonium chloride.....	2.54%
n-Akyl (C <sub>14</sub> 50%, C <sub>12</sub> 40%, C <sub>16</sub> 10%) dimethyl benzyl ammonium chloride.....	1.69%
Inert Ingredients:.....	95.77%
Total:.....	100.00%

## KEEP OUT OF REACH OF CHILDREN

# DA NGER

See side panel for Precautionary Statements

**FIRST AID**

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.  
**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.  
Call a poison control center or doctor for treatment advice.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.



Detco Industries  
P.O. Box 430 Conway, AR 72033  
www.detco.com

EPA Reg. No. 47371-131-58111  
EPA Est. No. 58111-AR-1

NET CONTENTS 1 Gal.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.  
**Pesticide Storage:** Open dumping is prohibited. Store in original container in areas inaccessible to children.  
**Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal or excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Fill container ¾ full with water and reclose the container. Agitate vigorously, and dispose of rinsate consistent with pesticide disposal instructions. Repeat two more times. Then offer for recycling if available or puncture and dispose in sanitary landfill or by other procedures approved by state and local authorities. Follow pesticide disposal instructions for rinsate. If not triple rinsed, these containers are acute hazardous wastes and must be disposed in accordance with local, state, and federal regulations.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER, Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Harmful if inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.**

Effective against SARS-CoV-2 in 1 minute.\*

Effective against the following pathogens:

Bacteria: <sup>1</sup>Pseudomonas aeruginosa, <sup>2</sup>Staphylococcus aureus<sup>1</sup>, Salmonella enterica, Acinetobacter baumannii, Acetobacter cacaeciarum, Bordetella bronchiseptica, Chlamydia psittaci, Enterobacter cloacae, Enterobacter cloacae - NDM-1, Enterococcus faecalis, Klebsiella aerogenes, Klebsiella pneumoniae<sup>1</sup>, Escherichia coli - NDM-1, Fusobacterium necrophorum, Klebsiella aerogenes, Klebsiella pneumoniae<sup>1</sup>, Klebsiella pneumoniae - NDM-1, Legionella pneumophila, Listeria monocytogenes, Pasteurella multocida, Proteus mirabilis, Proteus vulgaris, Salmonella enteritidis, Salmonella typhi, Serratia marcescens, Shigella flexneri, Shigella sonnei, Staphylococcus aureus - Community Associated Methicillin-Resistant (CA-MR-SAINRS394)(USA300), Staphylococcus aureus - Community Associated Methicillin-Resistant (CA-MRAS)(NRS123)(USA400), Staphylococcus aureus - Methicillin resistant - (MRSA), Staphylococcus aureus - Vancomycin Intermediate Resistant - (VISA), Staphylococcus epidermidis<sup>1</sup>, Streptococcus faecalis<sup>1</sup>, Streptococcus pyogenes (Strep)

Part # 0798-4X1  
Batch #



Viruses: <sup>3</sup>Adenovirus Type 7 - at 8 oz. per gallon, <sup>4</sup>Hepatitis B Virus (HBV), <sup>5</sup>Hepatitis C Virus (HCV), <sup>6</sup>Herpes Simplex Virus Type 1, <sup>7</sup>Herpes Simplex Virus Type 2, <sup>8</sup>Human coronavirus, <sup>9</sup>HIV-1 (AIDS virus), <sup>10</sup>Influenza A Virus, <sup>11</sup>Respiratory Syncytial Virus (RSV), <sup>12</sup>Rotavirus (WA), <sup>13</sup>Rubella virus, <sup>14</sup>SARS Associated Coronavirus (SARS), <sup>15</sup>SARS-Related Coronavirus 2(SARS-CoV-2), <sup>16</sup>Vaccinia virus

Animal Viruses: <sup>17</sup>Avian Influenza Virus H5N1, <sup>18</sup>Avian polyomavirus, <sup>19</sup>Canine distemper virus, <sup>20</sup>Feline leukemia virus, <sup>21</sup>Feline pneumovirus, <sup>22</sup>Infectious bovine rhinotracheitis virus, <sup>23</sup>Infectious bronchitis virus, <sup>24</sup>Newcastle Disease virus, <sup>25</sup>Pseudorabies virus, <sup>26</sup>Rabies virus, <sup>27</sup>Transmissible gastroenteritis virus

Fungi: <sup>28</sup>Aspergillus niger, <sup>29</sup>Candida albicans, <sup>30</sup>Trichophyton interdigitale (Athlete's Foot Fungus)

<sup>31</sup>ATCC & antibiotic-resistant strain, <sup>32</sup>antibiotic-resistant strain only, <sup>33</sup>Salmonella enterica, <sup>34</sup>hard, non-porous surfaces